

Neonatal circumcision: Risks and benefits

Author

Laurence S Baskin, MD,
FAAP

Section Editors

Charles J Lockwood, MD,
MHCM
John G Bartlett, MD
Duncan Wilcox, MD

Deputy Editor

Kristen Eckler, MD, FACOG

All topics are updated as new evidence becomes available and our [peer review process](#) is complete.

Literature review current through: May 2015. | **This topic last updated:** Apr 10, 2015.

INTRODUCTION — Circumcision in the male refers to the surgical removal of the prepuce (ie, foreskin) of the penis. The procedure is centuries old and continues to be performed for a variety of religious, cultural, and medical reasons.

Parents of newborn males often have questions about circumcision: Is it necessary? What are its benefits? What are its risks? In addition to these medical issues, other factors that influence parental decision-making include the father's circumcision status, opinions of family members and friends, a desire for conformity in their son's appearance, and the belief that the circumcised penis is easier to keep clean [1-3].

The risks and benefits of circumcision will be reviewed here, with the focus on circumcision of neonates. Procedures for neonatal circumcision are discussed separately. (See "[Techniques for neonatal circumcision](#)".)

BACKGROUND

Normal penile development and anatomy — (See "[Techniques for neonatal circumcision](#)", section on 'Normal development and anatomy'.)

Prevalence and epidemiology — The United States is the only country in the developed world where the majority of male infants are circumcised for nonreligious reasons. Circumcision rates in the United States vary according to geographic area, socioeconomic status, religious affiliation, insurance coverage, hospital type, and racial and ethnic group. The overall prevalence is estimated to be about 80 percent for males aged 14 to 59 years, with most of these procedures performed in newborns [4]. Circumcision rates are highest in the Midwestern states (74 percent), followed by the Northeastern states (67 percent) and Southern states (61 percent), and are lowest in the Western states (30 percent) [5]. The rate is higher in non-Hispanic whites (91 percent) than in non-Hispanic blacks (76 percent) and Mexican Americans (44 percent) [4].

Circumcision rates outside of the United States vary widely, from less than 5 percent to over 90 percent of males [6].

There are no studies that give reliable data about the number of males who are circumcised after birth for elective or surgical indications. An Australian survey found that approximately 18 percent of males who were not circumcised as infants reported that they were circumcised subsequently [7,8].

COUNSELING — Parents should receive factually correct, nonbiased information about the potential medical benefits and risks of neonatal circumcision and should understand that the procedure is elective. This discussion needs to consider the social, cultural, and religious aspects of the procedure, and the parents' personal preferences, which may be more important to them than a decision based on medical benefits alone [1-3,9,10]. In cross-sectional studies performed in the United States, 40 to 67 percent of parents reported they chose to have their son circumcised because of its hygienic and medical benefits; 23 to 37 percent had the procedure performed for social reasons (eg, father circumcised); and 11 to 19 percent of procedures were performed for religious reasons [11-14].

Ideally, counseling is provided when parents first begin to consider circumcision, which may be early in

pregnancy or before. Written informed consent is obtained prior to the procedure. (See "[Techniques for neonatal circumcision](#)", section on 'Informed consent'.)

Parents who are considering deferring circumcision until their son reaches an age where he can make an informed decision should be informed that circumcision performed in adults has greater risks and costs, more postprocedure discomfort, and a longer recovery period than newborn circumcision [15]. It also delays protection from sexually transmitted infections, which is important if the male becomes sexually active before the procedure is performed.

POTENTIAL BENEFITS — Circumcision has been associated with a number of medical benefits, including lower rates of urinary tract infection (UTI), penile cancer, penile inflammation, penile dermatoses, and sexually transmitted infections [16-20]. A lower rate of UTI is the major benefit during infancy.

In the uncircumcised male, the preputial space provides a warm moist environment that both traps pathogens and bodily secretions and is favorable to their survival and replication, while the foreskin itself is susceptible to microabrasions that are thought to facilitate acquisition and transmission of infection. In contrast, the keratinized glans of the circumcised male is thought to create an unfavorable environment for these processes.

Reduction in urinary tract infection — UTI is uncommon in males at any age. The effect of circumcision on UTI has been studied primarily in infants because they have a higher prevalence of UTI than older males. UTIs in infants can result in pyelonephritis requiring hospitalization and, rarely, septicemia and death. In infants with congenital uropathy, UTI can have serious sequelae, such as renal scarring and lifelong renal insufficiency. If the urinary tract is normal, long-term sequelae from UTI are unlikely.

Comparative studies consistently report that circumcised male infants have significantly fewer UTIs than uncircumcised male infants [17,18]; however, there are no data from a large randomized trial [21]. In a systematic review of 12 studies including data on over 400,000 males primarily under 1 year of age, circumcision reduced the risk of UTI by almost 90 percent (OR 0.13, 95% CI 0.08-0.20) [18]. Another meta-analysis found that among febrile male infants less than 3 months of age, the prevalence of UTI in circumcised and uncircumcised infants was 2.4 and 20.1 percent, respectively [17]. It is estimated that 7 to 14 of 1000 uncircumcised male infants will develop a UTI during the first year of life, compared with 1 to 2 of 1000 circumcised male infants [5]. As discussed above, long-term sequelae from UTI are unlikely if the urinary tract is normal.

There is some evidence that periurethral flora contains fewer pathogens after circumcision than before circumcision [22,23]. Uncircumcised male infants under 6 months of age have higher numbers of uropathogenic bacteria bound to the mucosal surface of the foreskin and at the urethral meatus than infants who have been circumcised [24]. This colonization probably plays a role in the pathogenesis of urinary infection. However, factors other than circumcision may affect the observed rate of UTI in these reports [25]. As an example, most hospitalized premature infants are not circumcised. Since premature infants have a higher rate of UTI than term infants, the inclusion of premature infants in a series may confound the data. Other factors that can affect UTI rates include the method of urine collection, type and timing of circumcision, and breastfeeding status (protective effect [26]).

The reduction in UTI associated with circumcision has also been observed after infancy. In a 2013 systematic review of primarily unrandomized studies, adult uncircumcised men had a higher rate of UTI than circumcised men (RR 3.4, 95% CI 0.92-12.7) and the lifetime risk of UTI in uncircumcised and circumcised men was 32.1 and 8.8 percent, respectively (RR 3.65, 95% CI 1.15-11.8) [27]. Lack of circumcision was estimated to account for 23 percent of UTIs during the lifetime of males.

The prevalence of UTIs in uncircumcised adult males increases with age and certain disease states, such as diabetes mellitus [28,29]. (See "[Acute uncomplicated cystitis, pyelonephritis, and asymptomatic bacteriuria in men](#)" and "[Urinary tract infections in children: Epidemiology and risk factors](#)", section on 'Lack of circumcision'.)

Infants with congenital uropathy — In male infants with serious congenital uropathies such as high grade

vesicoureteral reflux, posterior urethral valves, megaureters, and prune belly syndrome, reduction in urinary tract infections by circumcision may have greater potential benefit. Prospective randomized trials have not been performed, but the retrospective evidence described above supports this hypothesis. In my practice, I tell families that a circumcision has the potential to be beneficial, especially when the baby has an associated uropathy.

Reduction in risk of some cancers — Compared to uncircumcised men, circumcised men appear to have a lower risk of penile cancer, and their sexual partners may have a lower risk of cervical cancer.

Penile cancer — Squamous cell cancer of the penis is a rare disease. The age-adjusted incidence in the United States is less than 1 per 100,000 males, comparable to that in other developed countries [30,31].

A systematic review and meta-analysis found a strong protective effect of childhood/adolescent circumcision on invasive penile cancer (OR 0.33; 95% CI 0.13-0.83; 3 studies) [19]. In two studies, the protective effect did not persist when analyses were restricted to boys with no history of phimosis, suggesting at least some of the increased risk of cancer is related to phimosis, which can be the result of poor penile hygiene [32]. There was no association between circumcision and penile intraepithelial neoplasia and in situ penile cancer.

The number of newborn circumcisions that would need to be performed to prevent one case of penile cancer has been estimated to range from 909 to 322,000, and 2 to 644 complications would be expected per cancer prevented [33-35]. Given the low risk of penile cancer and the possibly modest reduction in risk with circumcision, it is unclear whether this benefit is clinically important when evaluated against the cost of the procedure and the risk of procedure related complications. (See "[Carcinoma of the penis: Epidemiology, risk factors, staging, and prognosis](#)", section on 'Risk factors'.) Furthermore, these data were obtained prior to widespread use of the human papillomavirus (HPV) vaccine for girls, as well as boys. (See "[Recommendations for the use of human papillomavirus vaccines](#)".) Vaccination may reduce the prevalence of HPV related disease, diminishing the role of circumcision in prevention of these diseases.

Cervical cancer in partners — Cervical cancer is more common in the sexual partners of uncircumcised men. A partial explanation for the link between cervical cancer and lack of male circumcision is that circumcised men have a lower prevalence of HPV infection than uncircumcised men [36], they are less likely to transmit HPV to their partners [37], and their partners have lower high-risk HPV DNA load [38]. (See "[Prevention of sexually transmitted infections](#)", section on 'Human papillomavirus'.)

In a study that pooled data from 1913 couples enrolled in one of seven case-control studies, circumcised men were less likely than uncircumcised men to have HPV infection (OR 0.37, 95% CI 0.16-0.85 after adjustment of confounders) and male circumcision was associated with a trend for reduced risk of cervical HPV and cervical cancer (OR 0.75, 95% CI 0.49-1.14) in the female partner [39]. Among monogamous women, the reduced risk of cervical cancer was statistically significant when the male partner had engaged in high-risk sexual behaviors (early age at first sexual intercourse, a high lifetime number of sexual partners, sexual intercourse with prostitutes) (OR 0.18, 95% CI 0.04-0.89) [39]. (See "[Cervical intraepithelial neoplasia: Terminology, incidence, pathogenesis, and prevention](#)", section on 'Role of human papillomavirus'.)

As discussed above, these data were obtained prior to widespread use of the HPV vaccine for girls, as well as boys. (See "[Recommendations for the use of human papillomavirus vaccines](#)".) Vaccination may reduce the prevalence of HPV related disease, diminishing the role of circumcision in prevention of these diseases.

Other — Case-control studies have reported that circumcised men have a lower rate of prostate cancer than uncircumcised men [40,41]. This could be related to multiple confounders; further investigation is required.

Reduction in penile inflammation and retractile disorders — In case series, penile problems are described more often in uncircumcised males [20,42,43]. The best available data were reported by a prospective cohort study of 635 New Zealand boys followed from birth to age 8 years [20]. The rates of penile inflammation and phimosis were higher in uncircumcised boys (penile inflammation: 14.4 versus 7.6 per 100 boys 0 to 8 years of age; phimosis 3.7 versus 0 per 100 boys 0 to 8 years of age).

Properly performed neonatal circumcision prevents phimosis, paraphimosis and balanoposthitis. A reduction in penile inflammatory and retractile disorders is important because, although they can often be treated medically [44-46], chronic inflammation and repeated forceful retraction may cause scarring and secondary phimosis, which sometimes requires surgical intervention [7,8,29,47]. However, good penile hygiene also may prevent penile inflammatory and retractile disorders, highlighting the importance of teaching uncircumcised boys routine penile care and hygiene. (See "[Care of the uncircumcised penis](#)" and "[Balanoposthitis in children: Clinical manifestations, diagnosis, and treatment](#)" and "[Paraphimosis: Definition, pathophysiology, and clinical features](#)".)

Reduction in HIV and other sexually transmitted infections — There is high-quality evidence that circumcision protects against acquisition of human immunodeficiency virus (HIV), HPV, and probably herpes simplex virus type 2 (HSV-2), and also some evidence that it may protect against trichomonas and chancroid infection (see "[Prevention of sexually transmitted infections](#)", section on 'Male circumcision'). Circumcision does not protect against infection from gonorrhea, chlamydia trachomatis, or syphilis.

The effects of male circumcision on preventing HIV transmission from men to women and on acquisition of other STIs are discussed elsewhere (See "[Prevention of sexually transmitted infections](#)", section on 'HIV infection in men' and "[Prevention of sexually transmitted infections](#)", section on 'Other sexually transmitted infections'.)

Easier hygiene — Good genital hygiene (washing the entire penis, including the glans, with soap and water while bathing) is important for all males and is generally easier in the absence of a foreskin.

Uncircumcised boys should be taught the importance of washing beneath the foreskin on a regular basis when the foreskin is fully retractable. Good hygiene may prevent many problems associated with the foreskin [32], but can be difficult to maintain in uncircumcised boys, even in developed countries. Studies of middle class British and Scandinavian schoolboys concluded that penile hygiene is usually not well-maintained [48,49]. Care of the uncircumcised penis is described separately. (See "[Care of the uncircumcised penis](#)".)

POTENTIAL DISADVANTAGES

Procedure related complications — An accurate complication rate is difficult to determine, as the largest studies are retrospective and based on coding diagnoses and inconsistent definitions. In addition, data have generally not been stratified to account for timing of the procedure, technique, provider type, setting, length of follow-up, timing of complications, and severity of complications.

- In two studies that included a total of over 200,000 circumcisions performed in US hospitals, the rate of complications during and in the first month after the procedure was approximately 0.2 percent [33,34].
- A systematic review identified 16 prospective studies of complications following neonatal and infant circumcision by a variety of providers from 12 countries and primarily using the Plastibell [50]. The median frequency of any adverse event was 1.5 percent (range 0 to 16 percent) and the median frequency of any serious adverse event was 0 percent (range 0 to 2 percent); nine studies reported no serious adverse events, but three studies reported that 1 to 2 percent of boys had a serious complication. Complication rates were slightly lower in 10 retrospective studies.

Complications/sequelae of circumcision are listed below and discussed in more detail separately (see "[Complications of circumcision](#)"):

- Inadequate skin removal, which may result in an unsatisfactory cosmetic appearance and revision of the procedure. This is a common complaint, although the frequency is poorly documented in the literature.
- Bleeding, which is usually mild and controlled with local pressure, but surgical intervention and transfusion may be required on rare occasions.
- Infection, which is usually mild and treated by local antibiotics, but sepsis can occur and death has been reported.

- Urethral complications, including urethrocutaneous fistula and meatal stenosis. Meatal stenosis can occur when urine from a wet diaper irritates the exposed ventral urethral meatus of the circumcised penis.
- Glans injury, including penile amputation
- Removal of excessive skin, which may result in a denuded penile shaft
- Epidermal inclusion cyst
- Adhesions, which range from mild to dense (also referred to as skin bridges)
- Cicatrix (a circumferential scar that usually develops at the incision line and is often associated with a hidden penis)
- Anesthetic complications (see ["Techniques for neonatal circumcision", section on 'Pain control'](#))

Complications are more common among premature newborns, newborns with congenital anomalies, and circumcisions performed after the newborn period or by untrained/poorly trained providers who may use nonsterile techniques [5].

Sexual dissatisfaction — The prepuce contains specialized sensory mucosa [51]; concern exists that the end of the penis may become less sensitive and sexual sensation may be decreased when the foreskin is removed [52]. However, most circumcised males do not describe psychological trauma or decreased sexual function, sensation, or desire as a result of the procedure. In a systemic review of 36 studies, circumcision was not associated with decreased sexual arousal, sensitivity, or satisfaction [53].

Pain — Surgical excision of the foreskin is painful. Although pain control was not provided in the past, safe and effective methods of pain control exist and should be provided to all infants undergoing the procedure [28,54]. The preferred method is subcutaneous ring or dorsal penile nerve block. Topical anesthetic creams are an acceptable alternative but should be avoided in infants who are low birth weight because of a higher incidence of skin irritation in this population. (See ["Techniques for neonatal circumcision", section on 'Pain control'](#).)

Other — In the era before use of anesthesia for circumcision, infants were observed to be less interactive and feed less frequently after the procedure [55,56]. This raised concerns that stress, pain, and fatigue related to circumcision may contribute to breastfeeding failure. However, changes in infant behavior do not appear to last longer than 24 hours [56], and anesthesia reduces procedure-related pain and its sequelae [57,58]. A longitudinal birth cohort study that examined the association between neonatal circumcision and breastfeeding found no evidence that neonatal circumcision disrupts exposure to, or outcomes of, breastfeeding [59].

OPINIONS OF ORGANIZATIONS AND GROUPS — Professional societies and lay groups have expressed a wide range of views concerning the advantages and disadvantages of routine circumcision. Several pediatric and urologic experts feel that circumcision should be advocated as a prophylactic public health measure [11,54,60-64], and there is increasing commentary from the pediatric literature suggesting that male circumcision may be an effective prophylactic intervention for disease prevention in the United States and other countries [65,66]. On the other hand, small, but vocal, groups have proposed the theory that all men who are circumcised may be emotionally harmed, but are unable to recognize their feelings. Other groups oppose circumcision on a human rights basis, saying the infant is not allowed to decide whether or not he wants to be circumcised and the procedure constitutes genital mutilation [62]. In 2012, a German court in Cologne ruled that circumcising young boys represents grievous bodily harm, but was overruled by a higher court. Still others believe the procedure decreases sexual pleasure [52].

- In 1996, the Canadian Paediatric Society (CPS) issued a clinical practice guideline which stated, "The overall evidence of the benefits and harms of circumcision is so evenly balanced that it does not support recommending circumcision as a routine procedure for newborns" [67].
- In 2012, the [American Academy of Pediatrics](#) (AAP) task force on circumcision of the male infant

concluded that "the health benefits of newborn male circumcision outweigh the risks; furthermore, the benefits of newborn male circumcision justify access to this procedure for families who choose it. Specific benefits from male circumcision were identified for the prevention of urinary tract infections, acquisition of HIV, transmission of some sexually transmitted infections, and penile cancer. Male circumcision does not appear to adversely affect penile sexual function/sensitivity or sexual satisfaction" [68]. Compared to their previous statement, this statement is a stronger affirmation of the health benefits of circumcision. However, the AAP did not recommend routine circumcision. They said "Parents should weigh the health benefits and risks in light of their own religious, cultural, and personal preferences, as the medical benefits alone may not outweigh these other considerations for individual families" [68].

- The American College of Obstetricians and Gynecologists (ACOG) endorsed the conclusions of the AAP discussed above [69].
- In 2012, the American Urological Association (AUA) reaffirmed their 2007 policy that stated, "The American Urological Association, Inc[®] (AUA) believes that neonatal circumcision has potential medical benefits and advantages as well as disadvantages and risks... is generally a safe procedure when performed by an experienced operator... When circumcision is being discussed with parents and informed consent obtained, medical benefits and risks, and ethnic, cultural, religious and individual preferences should be considered. The risks and disadvantages of circumcision are encountered early whereas the advantages and benefits are prospective" [70].

In addition, the World Health Organization (WHO) has recommended that male circumcision be considered as part of a comprehensive human immunodeficiency virus (HIV) prevention package in Africa and other countries with high HIV prevalence and low rates of male circumcision [71]. They stated that a human rights-based approach to promotion of male circumcision requires measures that ensure that the procedure be carried out safely, with informed consent, and without coercion or discrimination, and have published guidance for providing these services [72]. The effect of male circumcision on HIV transmission is discussed elsewhere. (See ["Prevention of sexually transmitted infections". section on 'Male circumcision'.](#))

INFORMATION FOR PATIENTS — UpToDate offers two types of patient education materials, "The Basics" and "Beyond the Basics." The Basics patient education pieces are written in plain language, at the 5th to 6th grade reading level, and they answer the four or five key questions a patient might have about a given condition. These articles are best for patients who want a general overview and who prefer short, easy-to-read materials. Beyond the Basics patient education pieces are longer, more sophisticated, and more detailed. These articles are written at the 10th to 12th grade reading level and are best for patients who want in-depth information and are comfortable with some medical jargon.

Here are the patient education articles that are relevant to this topic. We encourage you to print or e-mail these topics to your patients. (You can also locate patient education articles on a variety of subjects by searching on "patient info" and the keyword(s) of interest.)

- Basics topics (see ["Patient information: Circumcision in baby boys \(The Basics\)"](#) and ["Patient information: Caring for an uncircumcised penis \(The Basics\)"](#))
- Beyond the Basics topics (see ["Patient information: Circumcision in baby boys \(Beyond the Basics\)"](#))

SUMMARY AND RECOMMENDATIONS

- Male circumcision (removal of the prepuce or foreskin) is almost always an elective procedure performed for religious, cultural, or medical reasons. The prevalence varies across the world. (See ['Background'](#) above and ['Prevalence and epidemiology'](#) above.)
- Parents of newborn males should be provided with accurate, unbiased, written information about the potential risks and benefits of circumcision. This information should be in a language that they understand and they should have the opportunity to discuss this information with the operating practitioner prior to

deciding whether or not to have their infant son circumcised. (See '[Counseling](#)' above.)

- We suggest neonatal circumcision, as the health benefits of circumcision outweigh the risks (**Grade 2C**). Benefits of circumcision in men include reduction in the rates of urinary tract infection (UTI), penile cancer, some sexually transmitted infections, penile inflammatory and retractile disorders, as well as easier hygiene. These benefits, which extend over a lifetime, need to be weighed against the potential risks of the circumcision procedure, which are often short-term, and in the context of the low incidence of UTIs and penile cancer in uncircumcised men. (See '[Potential benefits](#)' above.)
- Circumcision protects the male from acquisition of human immunodeficiency virus (HIV), human papillomavirus (HPV), and probably herpes simplex virus type 2 (HSV-2). There is also some evidence that it may protect against trichomonas and chancroid infection. Male circumcision also confers benefits to female partners by reductions in cervical cancer and acquisition of some sexually transmitted infections and bacterial vaginosis. (See "[Prevention of sexually transmitted infections](#)", [section on 'Male circumcision'](#) and '[Cervical cancer in partners](#)' above.)
- Procedure-related complications are uncommon, and serious complications are rare. (See '[Procedure related complications](#)' above.)

Use of UpToDate is subject to the [Subscription and License Agreement](#).

REFERENCES

1. Neonatal circumcision. Report 10 of the Council on Scientific Affairs (I-99). www.ama-assn.org/ama/pub/category/13585.html (Accessed on September 14, 2006).
2. Waldeck SE. Social norm theory and male circumcision: why parents circumcise. *Am J Bioeth* 2003; 3:56.
3. Xu B, Goldman H. Newborn circumcision in Victoria, Australia: reasons and parental attitudes. *ANZ J Surg* 2008; 78:1019.
4. Morris BJ, Bailis SA, Wiswell TE. Circumcision rates in the United States: rising or falling? What effect might the new affirmative pediatric policy statement have? *Mayo Clin Proc* 2014; 89:677.
5. Blank S, Brady M, Buerk E, et al. Male circumcision. *Pediatrics* 2012; 130:e756.
6. Halperin DT, Timberg C. Update of chart in 'Male Circumcision and HIV Prevention: 10 Years and Counting' *Circs.org*. 2012. www.circs.org/library/HalperinTimberg (Accessed on December 14, 2006).
7. Badger, J. Circumcision. What you think. *Australian Forum* 1989; 2:10.
8. Badger, J. The great circumcision report part 2. *Australian Forum* 1989; 2:4.
9. Binner SL, Mastrobattista JM, Day MC, et al. Effect of parental education on decision-making about neonatal circumcision. *South Med J* 2002; 95:457.
10. Wang ML, Macklin EA, Tracy E, et al. Updated parental viewpoints on male neonatal circumcision in the United States. *Clin Pediatr (Phila)* 2010; 49:130.
11. Adler R, Ottaway MS, Gould S. Circumcision: we have heard from the experts; now let's hear from the parents. *Pediatrics* 2001; 107:E20.
12. Tiemstra JD. Factors affecting the circumcision decision. *J Am Board Fam Pract* 1999; 12:16.
13. Walton RE, Ostbye T, Campbell MK. Neonatal male circumcision after delisting in Ontario. Survey of new parents. *Can Fam Physician* 1997; 43:1241.
14. Turini GA 3rd, Reinert SE, McQuiston LD, Caldamone AA. Circumcision: a study of current parental decision-making. *Med Health R I* 2006; 89:365.
15. Morris BJ, Waskett JH, Banerjee J, et al. A 'snip' in time: what is the best age to circumcise? *BMC Pediatr* 2012; 12:20.
16. Mallon E, Hawkins D, Dinneen M, et al. Circumcision and genital dermatoses. *Arch Dermatol* 2000; 136:350.
17. Shaikh N, Morone NE, Bost JE, Farrell MH. Prevalence of urinary tract infection in childhood: a meta-

- analysis. *Pediatr Infect Dis J* 2008; 27:302.
18. Singh-Grewal D, Macdessi J, Craig J. Circumcision for the prevention of urinary tract infection in boys: a systematic review of randomised trials and observational studies. *Arch Dis Child* 2005; 90:853.
 19. Larke NL, Thomas SL, dos Santos Silva I, Weiss HA. Male circumcision and penile cancer: a systematic review and meta-analysis. *Cancer Causes Control* 2011; 22:1097.
 20. Fergusson DM, Lawton JM, Shannon FT. Neonatal circumcision and penile problems: an 8-year longitudinal study. *Pediatrics* 1988; 81:537.
 21. Jagannath VA, Fedorowicz Z, Sud V, et al. Routine neonatal circumcision for the prevention of urinary tract infections in infancy. *Cochrane Database Syst Rev* 2012; 11:CD009129.
 22. Günşar C, Kurutepe S, Alparslan O, et al. The effect of circumcision status on periurethral and glanular bacterial flora. *Urol Int* 2004; 72:212.
 23. Aridogan IA, Ilkit M, Izol V, et al. Glans penis and prepuce colonisation of yeast fungi in a paediatric population: pre- and postcircumcision results. *Mycoses* 2009; 52:49.
 24. Wiswell TE, Miller GM, Gelston HM Jr, et al. Effect of circumcision status on periurethral bacterial flora during the first year of life. *J Pediatr* 1988; 113:442.
 25. Circumcision policy statement. American Academy of Pediatrics. Task Force on Circumcision. *Pediatrics* 1999; 103:686.
 26. Pisacane A, Graziano L, Mazzarella G, et al. Breast-feeding and urinary tract infection. *J Pediatr* 1992; 120:87.
 27. Morris BJ, Wiswell TE. Circumcision and lifetime risk of urinary tract infection: a systematic review and meta-analysis. *J Urol* 2013; 189:2118.
 28. Spach DH, Stapleton AE, Stamm WE. Lack of circumcision increases the risk of urinary tract infection in young men. *JAMA* 1992; 267:679.
 29. Fakjian N, Hunter S, Cole GW, Miller J. An argument for circumcision. Prevention of balanitis in the adult. *Arch Dermatol* 1990; 126:1046.
 30. Barnholtz-Sloan JS, Maldonado JL, Pow-sang J, Giuliano AR. Incidence trends in primary malignant penile cancer. *Urol Oncol* 2007; 25:361.
 31. Frisch M, Friis S, Kjaer SK, Melbye M. Falling incidence of penis cancer in an uncircumcised population (Denmark 1943-90). *BMJ* 1995; 311:1471.
 32. Krueger H, Osborn L. Effects of hygiene among the uncircumcised. *J Fam Pract* 1986; 22:353.
 33. Wiswell TE, Geschke DW. Risks from circumcision during the first month of life compared with those for uncircumcised boys. *Pediatrics* 1989; 83:1011.
 34. Christakis DA, Harvey E, Zerr DM, et al. A trade-off analysis of routine newborn circumcision. *Pediatrics* 2000; 105:246.
 35. Learman LA. Neonatal circumcision: a dispassionate analysis. *Clin Obstet Gynecol* 1999; 42:849.
 36. Albero G, Castellsagué X, Giuliano AR, Bosch FX. Male circumcision and genital human papillomavirus: a systematic review and meta-analysis. *Sex Transm Dis* 2012; 39:104.
 37. Aynaud O, Piron D, Bijaoui G, Casanova JM. Developmental factors of urethral human papillomavirus lesions: correlation with circumcision. *BJU Int* 1999; 84:57.
 38. Davis MA, Gray RH, Grabowski MK, et al. Male circumcision decreases high-risk human papillomavirus viral load in female partners: a randomized trial in Rakai, Uganda. *Int J Cancer* 2013; 133:1247.
 39. Castellsagué X, Bosch FX, Muñoz N, et al. Male circumcision, penile human papillomavirus infection, and cervical cancer in female partners. *N Engl J Med* 2002; 346:1105.
 40. Ross RK, Shimizu H, Paganini-Hill A, et al. Case-control studies of prostate cancer in blacks and whites in southern California. *J Natl Cancer Inst* 1987; 78:869.
 41. Ewings P, Bowie C. A case-control study of cancer of the prostate in Somerset and east Devon. *Br J Cancer* 1996; 74:661.
 42. Herzog LW, Alvarez SR. The frequency of foreskin problems in uncircumcised children. *Am J Dis Child* 1986; 140:254.
 43. Kaplan GW. Complications of circumcision. *Urol Clin North Am* 1983; 10:543.

44. DeVries CR, Miller AK, Packer MG. Reduction of paraphimosis with hyaluronidase. *Urology* 1996; 48:464.
45. Wright JE. The treatment of childhood phimosis with topical steroid. *Aust N Z J Surg* 1994; 64:327.
46. Ashfield JE, Nickel KR, Siemens DR, et al. Treatment of phimosis with topical steroids in 194 children. *J Urol* 2003; 169:1106.
47. Stenram A, Malmfors G, Okmian L. Circumcision for phimosis--indications and results. *Acta Paediatr Scand* 1986; 75:321.
48. Oster J. Further fate of the foreskin. Incidence of preputial adhesions, phimosis, and smegma among Danish schoolboys. *Arch Dis Child* 1968; 43:200.
49. Kalcev, B. Circumcision and personal hygiene in school boys. *Med Officer* 1964; 122:171.
50. Weiss HA, Larke N, Halperin D, Schenker I. Complications of circumcision in male neonates, infants and children: a systematic review. *BMC Urol* 2010; 10:2.
51. Taylor JR, Lockwood AP, Taylor AJ. The prepuce: specialized mucosa of the penis and its loss to circumcision. *Br J Urol* 1996; 77:291.
52. Boyle GJ, Bensley GA. Adverse sexual and psychological effects of male infant circumcision. *Psychol Rep* 2001; 88:1105.
53. Morris BJ, Krieger JN. Does male circumcision affect sexual function, sensitivity, or satisfaction?--a systematic review. *J Sex Med* 2013; 10:2644.
54. Schoen EJ, Wiswell TE, Moses S. Reply to Carole M. Lannon and the task force on circumcision. *Pediatrics* 2001; 108:211.
55. Howard CR, Howard FM, Weitzman ML. Acetaminophen analgesia in neonatal circumcision: the effect on pain. *Pediatrics* 1994; 93:641.
56. Marshall RE, Porter FL, Rogers AG, et al. Circumcision: II. Effects upon mother-infant interaction. *Early Hum Dev* 1982; 7:367.
57. Brady-Fryer B, Wiebe N, Lander JA. Pain relief for neonatal circumcision. *Cochrane Database Syst Rev* 2004; :CD004217.
58. Dixon S, Snyder J, Holve R, Bromberger P. Behavioral effects of circumcision with and without anesthesia. *J Dev Behav Pediatr* 1984; 5:246.
59. Fergusson DM, Boden JM, Horwood LJ. Neonatal circumcision: effects on breastfeeding and outcomes associated with breastfeeding. *J Paediatr Child Health* 2008; 44:44.
60. Weiss GN, Weiss EB. A perspective on controversies over neonatal circumcision. *Clin Pediatr (Phila)* 1994; 33:726.
61. Schoen EJ. The status of circumcision of newborns. *N Engl J Med* 1990; 322:1308.
62. Poland RL. The question of routine neonatal circumcision. *N Engl J Med* 1990; 322:1312.
63. Schoen EJ, Wiswell TE, Moses S. New policy on circumcision--cause for concern. *Pediatrics* 2000; 105:620.
64. Weiss GN. Prophylactic neonatal surgery and infectious diseases. *Pediatr Infect Dis J* 1997; 16:727.
65. Flynn P, Havens P, Brady M, et al. Male circumcision for prevention of HIV and other sexually transmitted diseases. *Pediatrics* 2007; 119:821.
66. Dickerman JD, American Academy of Pediatrics. Circumcision in the time of HIV: when is there enough evidence to revise the American Academy of Pediatrics' policy on circumcision? *Pediatrics* 2007; 119:1006.
67. Neonatal circumcision revisited. Fetus and Newborn Committee, Canadian Paediatric Society. *CMAJ* 1996; 154:769.
68. American Academy of Pediatrics Task Force on Circumcision. Circumcision policy statement. *Pediatrics* 2012; 130:585.
69. The American College of Obstetricians and Gynecologists Endorses AAP Policy Statement on Circumcision. <http://www.aap.org/en-us/about-the-aap/aap-press-room/Pages/New-Benefits-Point-to-Greater-Benefits-of-Infant-Circumcision-But-Final-Say-is-Still-Up-to-parents-Says-AAP.aspx?nfstatus=401&nftoken=00000000-0000-0000-0000-000000000000&nfstatusdescription=ERROR%3a+No+local+token> (Accessed on August 28, 2012).

70. American Urological Association. Circumcision <http://www.auanet.org/content/aua-policies/policy-statements/c/circumcision.cfm> (Accessed on April 15, 2013).
71. UNAIDS. Safe, voluntary, informed male circumcision and comprehensive HIV prevention programming. March 2008. (Accessed December 15, 2008).
72. World Health Organization. Male circumcision quality assurance: a guide to enhancing the safety and quality of services. 2008.

Topic 6716 Version 26.0

Disclosures

Disclosures: **Laurence S Baskin, MD, FAAP** Nothing to disclose. **Charles J Lockwood, MD, MHCM** Consultant/Advisory Boards: Celula [Aneuploidy screening (Prenatal and cancer DNA screening tests in development)]. Equity Ownership/Stock Options: Celula [Aneuploidy screening (Prenatal and cancer DNA screening tests in development)]. **John G Bartlett, MD** Nothing to disclose. **Duncan Wilcox, MD** Nothing to disclose. **Kristen Eckler, MD, FACOG** Nothing to disclose.

Contributor disclosures are reviewed for conflicts of interest by the editorial group. When found, these are addressed by vetting through a multi-level review process, and through requirements for references to be provided to support the content. Appropriately referenced content is required of all authors and must conform to UpToDate standards of evidence.

Conflict of interest policy